MONDIAL POTATOES SPUNTA X VE66 295

TRADITIONAL FRESH

Maturity 50 Late Dormancy 60 Medium Yield mature 128 Very High Tuber size 83 Large Tuber shape Long Oval Number of tubers 12 - 14 Skin colour Light Yellow Flesh after cooking Light Yellow Cooking type B - Slightly mealy Dry matter content/Starch 19,7% / 13,8% UWW/Specific gravity 359 / 1 077 Internal bruising 12 Quite Sensitive Metribuzin sensitivity 70 Little Potato disorder 83


- Big size tubers
- Very attractive bright and smooth skin
- High yield
- Early tuberisation/maturing during short days
- Good storability













Resistances Foliage Blight 52 Tuber Blight 63 Alternaria Common scab Powdery scab Spraing PVY 15 Yntn tuber tolerance PCN Resistance Wart disease F1 Ro1/4 9 10 Ro2/3 *4 *8 F2 Pa2 3 F6 3 Pa3 F18








PRE-TREATMENT AND PLANTING

- A heat treatment is preferred to break the dormancy.
- Sprouts should be well hardened to avoid them breaking off.
- MONDIAL can be grown on all soil types.
- On light soils with free living nematodes there is a higher risk of spraying. Apply a nematicide on these soils.
- Can be planted with small white buds but good pre-sprouting can advance the crop.
- Early planting is advised.
- Prevent breaking of sprouts during planting.




GROWING ATTENTION POINTS

- Emergence is uniform with a good foliage development.
- Take care of a good soil moisture to prevent common scab.
- MONDIAL has a good drought tolerance.
- Preventive spraying against Phytophthora is advised.



HAULM KILLING AND HARVEST

- Use chemicals with a strong effect on stems to encourage easy tuber detachment.
- The skin should be fully set before harvest.
- MONDIAL is moderately susceptible to bruising. Reduce drop heights and avoid mechanical damage.
- Tubers are big. Harvest and store with care to avoid damage and bruising.



STORAGE

MONDIAL is suitable for long term storage

